

IFT-2004/30
 IPPP/04/85
 DCPT/04/170
 hep-ph/0412253
 February 2, 2008

CP violation in $B_d^0 \rightarrow \tau^+ \tau^-$ decays

Piotr H. Chankowski¹, Jan Kalinowski^{1,2}, Zbigniew Wąs³
 and Małgorzata Worek^{3,4}

¹ Institute of Theoretical Physics, Warsaw University, Hoża 69, 00-681, Warsaw, Poland

² IPPP, University of Durham, South Road, Durham DH1 3LE, UK

³ Institute of Nuclear Physics PAS, Radzikowskiego 152, 31-3420, Cracow, Poland

⁴ Institute of Nuclear Physics, NCSR "Demokritos", 15310 Athens, Greece

Abstract

Establishing CP violation in $B^0(\bar{B}^0) \rightarrow l^+ l^-$ decays requires a measurement of polarization of the final lepton pair, or a precise determination of the $B^0 \rightarrow l^+ l^-$ and $\bar{B}^0 \rightarrow l^+ l^-$ rates. We first argue that if the amplitudes of these decays are dominated by the scalar and pseudoscalar Higgs penguin diagrams, as happens *e.g.* in supersymmetry with large $\tan\beta$, the CP asymmetries depend practically on only one CP violating phase. This phase can be large, of the order of the CKM phase, leading to large CP asymmetries in the $\tau^+ \tau^-$ decay channel of $B_d^0(\bar{B}_d^0)$ mesons, potentially measurable in BELLE or BABAR experiments. Secondly, we show that the existing TAUOLA τ -lepton decay library supplemented by its universal interface can efficiently be used to search for $B^0(\bar{B}^0) \rightarrow \tau^+ \tau^-$ decays, and to investigate how the CP asymmetry is reflected in realistic experimental observables.

1 Introduction

40 years after its discovery in the Kaon system, CP violation has also been firmly established in B -physics in a series of high statistics experiments which enabled sufficiently precise measurements of the relevant observables. While the reported small difference between the time dependent CP asymmetries measured in $B \rightarrow J/\psi K_S$ and in $B \rightarrow \phi K_S$ decays [1], if confirmed, would conflict with the Standard Model (SM) prediction, there is as yet no convincing signal of a contribution to CP violation from physics beyond the SM. Such a signal may eventually be provided by yet more precise measurements and joint analysis of CP asymmetries measured in different channels. Equally important is to look for CP violation in channels in which no (or negligibly small) effects violating CP are predicted by the SM. Observation of a non-zero effect in such processes would be an unambiguous signal of new physics contribution to CP violation.

From many possible channels¹ we consider in this letter flavour changing decays of the neutral B mesons into lepton pairs, $B_{d,s}^0 \rightarrow l^+ l^-$. This channel has attracted a lot of attention since it is very sensitive to new physics which affects the b -quark Yukawa couplings [2–5]. Approximate and full one-loop calculations in the supersymmetric extension of the SM with large $\tan\beta$ (the ratio of the vacuum expectation values of the two Higgs doublets) [4–6] showed that in such a scenario one can expect truly spectacular enhancement of the rates of the decays $B_{d,s}^0 \rightarrow l^+ l^-$.

Moreover, new physics can also lead to observable CP violation in these decays, which is not predicted by the SM. For example, CP violation could manifest itself through non-equal leptonic decay rates of the tagged $B^0(t)$ and $\bar{B}^0(t)$ states ($B^0(t)$ and $\bar{B}^0(t)$ are the states which at $t = 0$ are tagged as B^0 and $\bar{B}^0$, respectively). If polarization of final state leptons can be determined, additional information on CP violation could be provided by non-equal $\Gamma(B^0(t) \rightarrow l_L^+ l_L^-)$ and $\Gamma(\bar{B}^0(t) \rightarrow l_R^+ l_R^-)$ [or $\Gamma(B^0(t) \rightarrow l_R^+ l_R^-)$ and $\Gamma(\bar{B}^0(t) \rightarrow l_L^+ l_L^-)$] decay rates [7, 8].

Theoretically leptonic decays are particularly clean as the only nonperturbative quantities, on which their rates depend, are the B^0 meson decay constants $F_{B_{d,s}}$. Moreover, $F_{B_{d,s}}$ cancel out in suitably defined CP asymmetries. On the other hand it is not clear whether the CP violation in these channels can be accessible experimentally. First of all these decays have not yet been seen: the best upper limit on the $B_d^0 \rightarrow \mu^+ \mu^-$ branching fraction comes at present from the BABAR experiment [9]:

$$Br(B_d^0 \rightarrow \mu^+ \mu^-) < 8.3 \times 10^{-8} , \quad (1)$$

which improves on previous limits of BELLE (1.6×10^{-7} [10]) and CDF (1.5×10^{-7} [11]). The $B_s^0 \rightarrow \mu^- \mu^+$ branching fraction is bounded by [12]

$$Br(B_s^0 \rightarrow \mu^+ \mu^-) < 2.7 \times 10^{-7} \quad (2)$$

resulting from the combination of the CDF (5.8×10^{-7} [11]) and D0 (4.1×10^{-7} [13]) (all limits are at 90% CL). The limits (1) and (2) are still about 2-3 orders of magnitude

¹For example, also very small CP violation is predicted by the SM in the charm sector.

above the corresponding predictions of the SM: $Br(B_d^0 \rightarrow \mu^+\mu^-) \approx 1.3 \times 10^{-10}$ and $Br(B_s^0 \rightarrow \mu^+\mu^-) \approx 3.6 \times 10^{-9}$ [14–16]. The main uncertainties of these predictions come from the decay constants F_{B_s} and F_{B_d} which are known up to a precision of $\sim 15\%$ [17]. If the decays $B_{d,s}^0 \rightarrow \mu^+\mu^-$ occur at the rates as predicted by the SM, their detection will become possible only at the LHC. New physics (like supersymmetry) can increase significantly their rates to a level that they can be observed at *BABAR*, *BELLE* or Tevatron in near future. However, as we will show, in this case the ratio of time integrated leptonic decay rates of the $B^0(t)$ and $\bar{B}^0(t)$ states to $\mu^+\mu^-$ is unlikely to deviate appreciably from unity. Polarization measurement also seems very difficult in the case of the $\mu^+\mu^-$ channel [18]. In the $\tau^+\tau^-$ channel the situation is quite different: large CP violating effects can be expected, and τ polarization measurement is possible. So far this channel has not been much explored experimentally on account of difficulties with identifying τ leptons. As a result, practically no experimental limits on $Br(B_{d,s}^0 \rightarrow \tau^+\tau^-)$ exist despite the fact that the corresponding rates are expected to be a factor $m_\tau^2/m_\mu^2 \sim 300$ larger than the ones for decays into $\mu^+\mu^-$ final states.

The purpose of this paper is twofold. Firstly, we point out that in realistic scenarios of large $\tan\beta$ MSSM, where the rates of $B^0(\bar{B}^0) \rightarrow l^+l^-$ decays are significantly increased to a level measurable at the running *BABAR* and *BELLE* experiments, the CP asymmetry in the $B_d^0(\bar{B}_d^0) \rightarrow \tau^+\tau^-$ channel can be quite large and potentially measurable. Secondly, we identify realistic experimental observables in which the CP asymmetry can be detected. In addition to the ratio of time integrated leptonic $B_d^0(t)$ and $\bar{B}_d^0(t)$ already mentioned, we consider two complementary observables: the $\pi^\pm$ energies from $\tau \rightarrow \pi\nu$ decays, and the acoplanarity angle between the decay planes of the ρ mesons which originate from $\tau \rightarrow \rho\nu$. The former is sensitive to the longitudinal, while the latter to the transverse polarizations of τ 's coming from B_d^0 and $\bar{B}_d^0$ decays. We show how the existing **TAUOLA** package [19–21] and its **universal interface** [22–24] can be used to search for these decays *and* for the CP violation, demonstrating that the necessary tools for full Monte Carlo simulations are reliable and ready for use in the $B_d^0(\bar{B}_d^0) \rightarrow \tau^+\tau^-$ physics.

2 General formulae

The effective Lagrangian describing $B_d^0(\bar{B}_d^0) \rightarrow l^+l^-$ decays can succinctly be written as

$$\mathcal{L}_{\text{eff}} = B_{s,d}^0 \bar{\psi}_l (b_{s,d} + a_{s,d} \gamma^5) \psi_l + \bar{B}_{s,d}^0 \bar{\psi}_l (\bar{b}_{s,d} + \bar{a}_{s,d} \gamma^5) \psi_l . \quad (3)$$

To simplify the notation the subscripts d and s referring to non-strange and strange B^0 mesons, unless explicitly written, will be omitted. Hermiticity (CPT invariance) implies

$$\bar{b} = b^* , \quad \bar{a} = -a^* . \quad (4)$$

The amplitudes of B^0 decays into two helicity eigenstates read

$$\begin{aligned} \mathcal{A}_L &\equiv \langle l_L^+ l_L^- | B_{s,d}^0 \rangle = M_B (a + b \beta) , \\ \mathcal{A}_R &\equiv \langle l_R^+ l_R^- | B_{s,d}^0 \rangle = M_B (a - b \beta) , \end{aligned} \quad (5)$$

where $\beta = (1 - 4m_l^2/M_B^2)^{1/2}$. Similar formulae with a and b replaced by $\bar{a}$ and $\bar{b}$, respectively, give the amplitudes $\bar{\mathcal{A}}_L$ and $\bar{\mathcal{A}}_R$ for the corresponding $\bar{B}^0$ decays. Evidently, if both coefficients a and b are simultaneously nonzero, $\Gamma(B^0 \rightarrow l_L^+ l_L^-) \neq \Gamma(B^0 \rightarrow l_R^+ l_R^-)$ but this is not yet a signal of CP violation. CP is violated, for example, if

$$\Gamma(B^0 \rightarrow l_L^+ l_L^-) \neq \Gamma(\bar{B}^0 \rightarrow l_R^+ l_R^-) , \quad (6)$$

$$\Gamma(B^0 \rightarrow l_R^+ l_R^-) \neq \Gamma(\bar{B}^0 \rightarrow l_L^+ l_L^-) , \quad (7)$$

because the initial and final states on both sides transform into each other under CP [7]. As there are no strong phases involved, this can occur only through the mixing of the B^0 and $\bar{B}^0$ mesons. In the standard formalism [25] the state $B_{\text{phys}}^0(t)$ ($\bar{B}_{\text{phys}}^0(t)$) which at $t = 0$ is a pure B^0 ($\bar{B}^0$) evolves in time according to

$$\begin{aligned} |B_{\text{phys}}^0(t)\rangle &= g_+(t)|B^0\rangle + \frac{q}{p}g_-(t)|\bar{B}^0\rangle , \\ |\bar{B}_{\text{phys}}^0(t)\rangle &= g_+(t)|\bar{B}^0\rangle + \frac{p}{q}g_-(t)|B^0\rangle . \end{aligned}$$

Neglecting the difference of the decay widths of the two B^0 mass eigenstates one finds

$$\begin{aligned} g_+(t) &= e^{-iM_B t - \frac{\Gamma}{2}t} \cos \frac{\Delta M}{2}t , \\ g_-(t) &= e^{-iM_B t - \frac{\Gamma}{2}t} i \sin \frac{\Delta M}{2}t , \end{aligned} \quad (8)$$

where $\Delta M \equiv M_{B_H^0} - M_{B_L^0} \ll M_B \equiv (M_{B_H^0} + M_{B_L^0})/2$. The ratio p/q (calculable in the SM or its extensions) is given by

$$\frac{p}{q} = \sqrt{\frac{H_{12}^*}{H_{12}}} \approx \frac{M_{12}^*}{|M_{12}|} \left(1 - \frac{1}{2} \text{Im} \frac{\Gamma_{12}}{M_{12}}\right) , \quad (9)$$

with $H_{12} \equiv M_{12} + \frac{i}{2}\Gamma_{12} = \langle B^0 | \mathcal{H}_{\text{eff}} | \bar{B}^0 \rangle$, *etc.* [25]. The probability amplitude that the state, which initially was a B^0 , decays at time t into left-handed leptons is therefore given by

$$\langle l_L^+ l_L^- | B_{\text{phys}}^0(t) \rangle = g_+(t) \mathcal{A}_L + \frac{q}{p} g_-(t) \bar{\mathcal{A}}_L , \quad (10)$$

and the rates of $B^0 \rightarrow l_L^+ l_L^-$ and $\bar{B}^0 \rightarrow l_R^+ l_R^-$ decays are proportional to

$$\begin{aligned} |\langle l_L^+ l_L^- | B_{\text{phys}}^0(t) \rangle|^2 &= |g_+(t)|^2 |\mathcal{A}_L|^2 \left\{ 1 + \left| \frac{q g_-(t)}{p g_+(t)} \right|^2 \left| \frac{\bar{\mathcal{A}}_L}{\mathcal{A}_L} \right|^2 + 2 \text{Re} \left(\frac{q g_-(t)}{p g_+(t)} \frac{\bar{\mathcal{A}}_L}{\mathcal{A}_L} \right) \right\} , \\ |\langle l_R^+ l_R^- | \bar{B}_{\text{phys}}^0(t) \rangle|^2 &= |g_+(t)|^2 |\bar{\mathcal{A}}_R|^2 \left\{ 1 + \left| \frac{p g_-(t)}{q g_+(t)} \right|^2 \left| \frac{\mathcal{A}_R}{\bar{\mathcal{A}}_R} \right|^2 + 2 \text{Re} \left(\frac{p g_-(t)}{q g_+(t)} \frac{\mathcal{A}_R}{\bar{\mathcal{A}}_R} \right) \right\} . \end{aligned}$$

The matrix elements for $B^0 \rightarrow l_R^+ l_R^-$ and $\bar{B}^0 \rightarrow l_L^+ l_L^-$ can be obtained from the ones above by interchanging $L \leftrightarrow R$.

Since $|\mathcal{A}_L| = |\bar{\mathcal{A}}_R|$, $|\mathcal{A}_R| = |\bar{\mathcal{A}}_L|$ and, as follows from (4) and (5),

$$\frac{\bar{\mathcal{A}}_L}{\mathcal{A}_L} = \left(\frac{\mathcal{A}_R}{\bar{\mathcal{A}}_R} \right)^* , \quad (11)$$

one sees that the CP is violated if either

$$\left| \frac{q}{p} \right| \neq 1 \quad \text{or} \quad \text{Im}(\lambda_L) \neq \text{Im}(\lambda_R^{-1}) ,$$

where

$$\lambda_L \equiv \frac{q}{p} \frac{\bar{\mathcal{A}}_L}{\mathcal{A}_L} , \quad \text{and} \quad \lambda_R \equiv \frac{q}{p} \frac{\bar{\mathcal{A}}_R}{\mathcal{A}_R} . \quad (12)$$

The simplest quantities measuring the amount of CP violation are the asymmetries constructed out of time integrated polarized decay rates

$$A_{\text{CP}}^1(t_1, t_2) \equiv \frac{\int_{t_1}^{t_2} dt \Gamma(B_{\text{phys}}^0(t) \rightarrow l_L^+ l_L^-) - \int_{t_1}^{t_2} dt \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow l_R^+ l_R^-)}{\int_{t_1}^{t_2} dt \Gamma(B_{\text{phys}}^0(t) \rightarrow l_L^+ l_L^-) + \int_{t_1}^{t_2} dt \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow l_R^+ l_R^-)} , \quad (13)$$

$$A_{\text{CP}}^2(t_1, t_2) \equiv \frac{\int_{t_1}^{t_2} dt \Gamma(B_{\text{phys}}^0(t) \rightarrow l_R^+ l_R^-) - \int_{t_1}^{t_2} dt \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow l_L^+ l_L^-)}{\int_{t_1}^{t_2} dt \Gamma(B_{\text{phys}}^0(t) \rightarrow l_R^+ l_R^-) + \int_{t_1}^{t_2} dt \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow l_L^+ l_L^-)} . \quad (14)$$

and the ratio of integrated unpolarized decay rates

$$R_l(t_1, t_2) \equiv \frac{\int_{t_1}^{t_2} dt \Gamma(B_{\text{phys}}^0(t) \rightarrow l^+ l^-)}{\int_{t_1}^{t_2} dt \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow l^+ l^-)} . \quad (15)$$

The time interval (t_1, t_2) can be chosen according to the experimental convenience. If the number of tagged events is not very large, or there is a large uncertainty in experimental determination of the decay time t , one can exploit the asymmetries $A_{\text{CP}}^1 \equiv A_{\text{CP}}^1(0, \infty)$ and $A_{\text{CP}}^2 \equiv A_{\text{CP}}^2(0, \infty)$ and the ratio $R_l \equiv R_l(0, \infty)$ for which it is straightforward to obtain [7, 26]

$$A_{\text{CP}}^1 = \frac{\frac{1}{2}x^2 (|\lambda_L^2| - |\lambda_R^{-2}|) - x \text{Im}(\lambda_L - \lambda_R^{-1})}{2 + x^2 + \frac{1}{2}x^2 (|\lambda_L^2| + |\lambda_R^{-2}|) - x \text{Im}(\lambda_L + \lambda_R^{-1})} , \quad (16)$$

$$A_{\text{CP}}^2 = \frac{\frac{1}{2}x^2 (|\lambda_R^2| - |\lambda_L^{-2}|) - x \text{Im}(\lambda_R - \lambda_L^{-1})}{2 + x^2 + \frac{1}{2}x^2 (|\lambda_R^2| + |\lambda_L^{-2}|) - x \text{Im}(\lambda_R + \lambda_L^{-1})} , \quad (17)$$

and

$$R_l = \frac{(|\mathcal{A}_L|^2 + |\mathcal{A}_R|^2) (1 + \frac{1}{2}x^2 + \frac{1}{2}x^2 |\frac{q}{p}|^2) - x \{ |\mathcal{A}_L|^2 \text{Im}(\lambda_L) + |\mathcal{A}_R|^2 \text{Im}(\lambda_R) \}}{(|\mathcal{A}_L|^2 + |\mathcal{A}_R|^2) (1 + \frac{1}{2}x^2 + \frac{1}{2}x^2 |\frac{p}{q}|^2) - x \{ |\mathcal{A}_L|^2 \text{Im}(\lambda_R^{-1}) + |\mathcal{A}_R|^2 \text{Im}(\lambda_L^{-1}) \}} , \quad (18)$$

where $x \equiv \Delta M/\Gamma$. If $|q/p| = 1$, the relation (11) implies $|\lambda_L| = |\lambda_R^{-1}|$. Moreover, $\lambda_L + \lambda_R^{-1}$ is then real, and the formulae (16), (17) and (18) simplify to

$$\begin{aligned} A_{\text{CP}}^1 &= \frac{-2x \operatorname{Im} \lambda_L}{2 + x^2 + x^2 |\lambda_L|^2}, & A_{\text{CP}}^2 &= \frac{-2x \operatorname{Im} \lambda_R}{2 + x^2 + x^2 |\lambda_R|^2}, \\ R_l &= \frac{(|\mathcal{A}_L|^2 + |\mathcal{A}_R|^2)(1 + x^2) - x \{|\mathcal{A}_L|^2 \operatorname{Im}(\lambda_L) + |\mathcal{A}_R|^2 \operatorname{Im}(\lambda_R)\}}{(|\mathcal{A}_L|^2 + |\mathcal{A}_R|^2)(1 + x^2) + x \{|\mathcal{A}_L|^2 \operatorname{Im}(\lambda_L) + |\mathcal{A}_R|^2 \operatorname{Im}(\lambda_R)\}}. \end{aligned} \quad (19)$$

The asymmetries A_{CP}^1 , A_{CP}^2 , as functions of $\lambda_{L,R}$, are bounded from above by [7]

$$|A_{\text{CP}}^{1,2}| \leq \frac{1}{\sqrt{2 + x^2}}. \quad (20)$$

Since $x_s > 20.6$ for the B_s^0 - $\bar{B}_s^0$ system, the CP asymmetries in the leptonic $B_s^0(\bar{B}_s^0)$ decays can reach at best $\sim 4.5\%$. In contrast, for the B_d^0 - $\bar{B}_d^0$ system, for which $x_d = 0.771 \pm 0.012$ they can be as large as $\sim 60\%$. This is fortunate, since $B_d^0 \bar{B}_d^0$ are copiously produced at $B\text{ABAR}$ and BELLE in a relatively clean environment (compared to the B^0 production at hadron colliders). For this reason we will consider only the CP asymmetries in the $B_d^0(\bar{B}_d^0) \rightarrow l^+ l^-$ decays.

The quantities (13) and (14) depend on asymmetries of B^0 and $\bar{B}^0$ decays into longitudinally polarized leptons. In the case of the $\tau^+ \tau^-$ decay mode they are best identified by measuring the $\pi^\pm$ energy spectra from $\tau \rightarrow \pi \nu$ decays [27]. The density matrix formalism outlined in Section 4 allows to construct also observables sensitive to transverse polarization of the final state τ 's [28]. These observables will prove complementary since in some scenarios the signal of CP violation can clearly be visible in the latter observables while hidden in the former (as R_l can be expressed in terms of polarized decay rates, R_l is then equal 1).

3 Supersymmetry scenario

If the Cabibbo-Kobayashi-Maskawa (CKM) matrix is the only source of *both* flavour and CP violation (as in the SM), then $q/p \approx -V_{tb}^* V_{td(s)}/V_{tb} V_{td(s)}^*$ implying $|q/p| \approx 1$. Moreover, $\mathcal{A}_L, \mathcal{A}_R \propto V_{tb}^* V_{td(s)}$, $\bar{\mathcal{A}}_L, \bar{\mathcal{A}}_R \propto V_{tb} V_{td(s)}^*$, so that λ_L and λ_R are almost real. The CP asymmetries in the $B^0 \rightarrow l^+ l^-$ decay are then negligible.² In models of new physics one can have $\operatorname{Im}(\lambda_L) \neq \operatorname{Im}(\lambda_R^{-1})$ and/or $|q/p| \neq 1$, but if the predicted rates of these decays are still of the same order as the SM predicts, the detection of the $\mu^+ \mu^-$ decay mode will become possible only at the LHC where most probably the measurement of the muon polarization will not be feasible. Detecting CP violation in this mode could be possible then only by measuring the ratio R_μ . The detection of the $B_d^0 \rightarrow \tau^+ \tau^-$ mode occurring at the SM model level ($Br \sim 5 \times 10^{-8}$) is possible at BELLE and $B\text{ABAR}$ ³ but the number of

²In the SM they are estimated to be $\mathcal{O}(10^{-2})$ [7] as a result of a small departure of $|q/p|$ from 1.

³According to the SM prediction, among 5×10^8 events of $B_d^0 \bar{B}_d^0$ pair production collected by these experiments so far there should already be some 50 events of $B_d^0(\bar{B}_d^0)$ decays into $\tau^+ \tau^-$.

reconstructed events might be too small to detect any CP violation. On the other hand, at LHC where the number of $B_{d,s}^0 \rightarrow \tau^+ \tau^-$ events will be larger, the identification of the τ decay mode will probably be quite difficult and the CP asymmetry hard to detect.

Much more promising situation can occur in the supersymmetric scenario with a large ratio of the vacuum expectation values of the two Higgs doublets, $v_u/v_d \equiv \tan \beta \sim 40 \div 50$. The coefficients a and b (and $\bar{a}$ and $\bar{b}$) in (3), and hence, the amplitudes of the $B_{d,s}^0 \rightarrow l^+ l^-$ decays can then receive important contributions from the Higgs penguin diagrams with s -channel H^0 and A^0 Higgs boson exchanges [4, 5, 8]. If the mass scale of the Higgs particles H^0 and A^0 (for $v_u/v_d \gg 1$ $M_H \approx M_A$) is not too high, of order $\lesssim 500$ GeV, the decay amplitudes can be dominated totally by these diagrams easily saturating the experimental limits (1), (2). This can happen even if the supersymmetric particle masses are quite large, say in the TeV range [5, 29].

As illustrative examples of new physics we consider here two different supersymmetric scenarios: minimal (MFV) and non-minimal (NMFV) flavour violating, both with large ratio of VEVs.

MFV Scenario: In this case no additional flavour violation in the sfermion mass matrices is assumed (*i.e.* the gluino-quark-squark vertices conserve flavour) but additional CP violation phases are introduced by complex parameters μ and A_t (the higgsino mass term and left-right top squark mixing, respectively). The relevant effective flavour violating couplings of the two heavy neutral Higgs bosons to the down-type quarks⁴ can be written as

$$\begin{aligned} \mathcal{L}_{\text{eff}}^{H^0, A^0 bd} = & C V_{tb}^* V_{td} \bar{b} \left[\frac{m_b}{M_W} \frac{A_t^*}{\mu} (H^0 + iA^0) \mathbf{P}_L + \frac{m_d}{M_W} \frac{A_t}{\mu^*} (H^0 - iA^0) \mathbf{P}_R \right] d \\ & + C V_{td}^* V_{tb} \bar{d} \left[\frac{m_d}{M_W} \frac{A_t^*}{\mu} (H^0 + iA^0) \mathbf{P}_L + \frac{m_b}{M_W} \frac{A_t}{\mu^*} (H^0 - iA^0) \mathbf{P}_R \right] b, \end{aligned}$$

where $\mathbf{P}_{L,R} = (1 \mp \gamma^5)/2$ are chiral projectors and the coefficient C is given by

$$C = \frac{g^3}{4} \frac{m_t^2}{M_W^2} \tan^2 \beta \frac{H_2}{16\pi^2} \kappa$$

with a dimensionless function of higgsino and stop masses H_2 of order $\mathcal{O}(1)$. The factor $\kappa \propto \tan \beta$ summarises some refinements in the calculation (resummation of $\tan \beta$ enhanced terms, for details see [6, 29]). It is the factor $\tan^2 \beta$ which makes these couplings so important. Combining these couplings with the H^0 and A^0 couplings to $l^+ l^-$ and using

$$\begin{aligned} \langle 0 | \bar{b} \mathbf{P}_L d | B_d^0 \rangle &= -\frac{i}{2} F_{B_d} \frac{M_B^2}{m_b}, & \langle 0 | \bar{d} \mathbf{P}_L b | \bar{B}_d^0 \rangle &= +\frac{i}{2} F_{B_d} \frac{M_B^2}{m_b}, \\ \langle 0 | \bar{b} \mathbf{P}_R d | B_d^0 \rangle &= +\frac{i}{2} F_{B_d} \frac{M_B^2}{m_b}, & \langle 0 | \bar{d} \mathbf{P}_R b | \bar{B}_d^0 \rangle &= -\frac{i}{2} F_{B_d} \frac{M_B^2}{m_b}, \end{aligned}$$

⁴The effective flavour violating coupling of the lightest CP even Higgs h^0 to the down-type quarks is negligible.

one arrives at

$$a = -\bar{a}^* = C' V_{tb}^* V_{td} \frac{M_B^2}{M_A^2} \left(\frac{m_b}{M_W} \frac{A_t^*}{\mu} + \frac{m_d}{M_W} \frac{A_t}{\mu^*} \right) \approx C' V_{tb}^* V_{td} \frac{m_b}{M_W} \frac{M_B^2}{M_A^2} \frac{A_t^*}{\mu} ,$$

$$b = \bar{b}^* = C' V_{tb}^* V_{td} \frac{M_B^2}{M_H^2} \left(\frac{m_b}{M_W} \frac{A_t^*}{\mu} - \frac{m_d}{M_W} \frac{A_t}{\mu^*} \right) \approx C' V_{tb}^* V_{td} \frac{m_b}{M_W} \frac{M_B^2}{M_H^2} \frac{A_t^*}{\mu} ,$$

with

$$C' = -\frac{g^4}{16} \frac{m_t^2}{M_W^2} \frac{m_l}{M_W} \frac{F_B}{m_b} \tan^3 \beta \frac{H_2}{16\pi^2} \kappa$$

In this case the amplitude of the B_d^0 - $\bar{B}_d^0$ mixing is not modified (in contrast to the one for B_s^0 - $\bar{B}_s^0$ mixing) [29], so that $|q/p| \approx 1$ holds true as in the SM. One then finds

$$\lambda_L \approx e^{-2i\delta_{\text{CP}}} \frac{1-\beta}{1+\beta} , \quad \lambda_R \approx e^{-2i\delta_{\text{CP}}} \frac{1+\beta}{1-\beta} , \quad (21)$$

where the effective CP violating phase is given as $\delta_{\text{CP}} = \arg(\mu^* A_t^*)$. The time integrated asymmetries then read

$$A_{\text{CP}}^1 = -0.09 \times \sin(2\delta_{\text{CP}}) , \quad A_{\text{CP}}^2 = -0.35 \times \sin(2\delta_{\text{CP}}) . \quad (22)$$

Since the sparticles giving rise to substantial Higgs penguin contributions to a and b ($\bar{a}$ and $\bar{b}$) can be quite heavy, even order $\mathcal{O}(1)$ phase of μA_t needs not produce unacceptable electric dipole moments.

NMFV Scenario: In this case squark mass matrices violate flavour and the corrections generating the flavour changing couplings of the neutral Higgs bosons H^0 and A^0 are dominated by gluino loops [5]. The relevant effective Lagrangian is then

$$\mathcal{L}_{\text{eff}}^{H^0, A^0 bd} = D \bar{b} \left[\alpha_b (H^0 + iA^0) \mathbf{P}_L + \alpha_d^* (H^0 - iA^0) \mathbf{P}_R \right] d$$

$$+ D \bar{d} \left[\alpha_d (H^0 + iA^0) \mathbf{P}_L + \alpha_b^* (H^0 - iA^0) \mathbf{P}_R \right] b , \quad (23)$$

with

$$\alpha_b = \frac{m_{\tilde{g}}^*}{\mu} \left(\frac{m_b}{M_W} \delta_{LL}^{bd} + \frac{m_d}{M_W} \delta_{RR}^{bd} \right) , \quad \alpha_d = \frac{m_{\tilde{g}}^*}{\mu} \left(\frac{m_d}{M_W} \delta_{LL}^{db} + \frac{m_b}{M_W} \delta_{RR}^{db} \right)$$

and

$$D = \frac{4}{3} g_s^2 g \tan^2 \beta \frac{|\mu|^2}{M_{\tilde{q}}^2} \frac{H_3}{16\pi^2} \kappa ,$$

where H_3 is another dimensionless function of order $\mathcal{O}(1)$ of gluino and sbottom masses and κ again denotes dominant higher order contributions [30]. The mass insertions δ_{LL}^{bd}

and δ_{RR}^{bd} etc. are the ratios of the diagonal entries of the down-type squark mass squared matrices to the average squark mass squared $M_{\tilde{q}}^2$. One then finds

$$a = -\bar{a}^* = D' (\alpha_b + \alpha_d^*) \approx D' \frac{m_b}{M_W} \frac{M_B^2}{M_A^2} \left(\frac{m_{\tilde{g}}^*}{\mu} \delta_{LL}^{bd} + \frac{m_{\tilde{g}}}{\mu^*} \delta_{RR}^{db*} \right) ,$$

$$b = \bar{b}^* = D' (\alpha_b - \alpha_d^*) \approx D' \frac{m_b}{M_W} \frac{M_B^2}{M_H^2} \left(\frac{m_{\tilde{g}}^*}{\mu} \delta_{LL}^{bd} - \frac{m_{\tilde{g}}}{\mu^*} \delta_{RR}^{db*} \right) ,$$

with

$$D' = -\frac{1}{3} g_s^2 g^2 \frac{m_l}{M_W} \tan^3 \beta \frac{|\mu|^2}{M_{\tilde{q}}^2} \frac{H_3}{16\pi^2} \kappa \frac{F_B}{m_b} .$$

The insertions δ_{LL}^{bd} and δ_{RR}^{bd} are not very tightly constrained. Typically the bound of order $|\delta_{LL(RR)}^{bd}| < 0.2 \times (M_{\tilde{q}}/1 \text{ TeV})$ arising from ΔM_{B_d} is quoted [31]. This estimate, however, does not take into account the contribution of the so-called double-penguin diagrams [30, 32–34], which can significantly affect the B_d^0 - $\bar{B}_d^0$ mixing amplitude. This contribution arises from the H^0 and A^0 exchanges between two effective (1-loop generated) vertices (23) of which one vertex annihilates a right-chiral and the other a left-chiral quark⁵ and is therefore proportional to $\alpha_b \alpha_d^* \propto \delta_{LL}^{bd} \delta_{RR}^{db*} = \delta_{LL}^{bd} \delta_{RR}^{bd}$ (with the small d-quark mass neglected). Since this product is constrained much stronger [30], to avoid a potential conflict with the value of ΔM_{B_d} and the time dependent CP asymmetry $a_{J/\psi K_S}(t)$ measured in $B \rightarrow J/\psi K_S$ decay, we assume that either $\delta_{RR}^{bd} \gg \delta_{LL}^{bd}$ or $\delta_{RR}^{bd} \ll \delta_{LL}^{bd}$. This leads to $a \approx \pm b$ and, as in the previous scenario, to $|q/p| \approx 1$. The asymmetries then read

$$A_{\text{CP}}^1 = -0.35 \times \sin(2\delta_{\text{CP}}) , \quad A_{\text{CP}}^2 = -0.09 \times \sin(2\delta_{\text{CP}}) ,$$

where now

$$\delta_{\text{CP}} = \arg(V_{tb}^* V_{td}) - \arg(m_{\tilde{g}} \mu \delta_{LL(RR)}^{bd}) . \quad (24)$$

It is interesting to note here that the CP asymmetries can be nonzero even if the phase of the CKM matrix remains the only source of CP violation (*i.e.* all supersymmetric parameters are real). Moreover, since $|\arg(V_{tb}^* V_{td})|$ is of order 1, the total phase violating CP needs not be small.

In both scenarios, in which the $B_{d,s}^0 \rightarrow l^+ l^-$ amplitudes are dominated by the exchange of H^0 and A^0 Higgs bosons whose effective flavour violating couplings to bd (or bs) differ only by a factor i , one gets

$$a \approx b \quad \text{or} \quad a \approx -b ,$$

⁵Contributions of H^0 and A^0 exchanges between vertices annihilating same chirality quarks to the mixing amplitude are proportional to $1/M_H^2 - 1/M_A^2 \approx 0$ [32].

(up to $\lesssim 15\%$). For $a = b$ the factors λ_L and λ_R defined in Eq. (12) simplify to:

$$\lambda_L = -\frac{q}{p} \frac{a^*}{a} \frac{1 - \beta}{1 + \beta}, \quad \lambda_R = -\frac{q}{p} \frac{a^*}{a} \frac{1 + \beta}{1 - \beta} \quad (25)$$

and all CP-sensitive quantities depend on one effective phase which can be taken as

$$\delta_{\text{CP}} = -\frac{1}{2} \arg(\lambda_L). \quad (26)$$

The immediate consequence of $a = b$, with $|q/p|$ close to 1, is that for the $\mu^+\mu^-$ final state the parameters $|\lambda_L|$ and $|\lambda_R|$ assume values $\sim 4 \times 10^{-4}$ and $\sim 2.5 \times 10^3$, respectively, since in this case $\beta = (1 - 4m_\mu^2/M_B^2)^{1/2}$ is almost 1. As a result, the predicted asymmetries are very small: $|A_{\text{CP}}^1| \lesssim 2 \times 10^{-4}$, $|A_{\text{CP}}^2| \lesssim 10^{-3}$. The same is true if a and b are somewhat split. In contrast, for the $\tau^+\tau^-$ final states, for which $\beta = (1 - 4m_\tau^2/M_B^2)^{1/2}$ differs substantially from 1, we have $|\lambda_L| \sim 0.15$, $|\lambda_R| \sim 6.7$ and the maximal possible values of the asymmetries are

$$|(A_{\text{CP}}^1)^{\text{max}}| = 9\% \quad \text{and} \quad |(A_{\text{CP}}^2)^{\text{max}}| = 35\%. \quad (27)$$

The comparison of magnitudes of possible CP violating effects in the ratio (18) for $\mu^+\mu^-$ and $\tau^+\tau^-$ decay modes is shown in Fig. 1, where R_μ and R_τ are plotted as functions of b/a for four different values of the phase δ_{CP} (keeping $\arg(a) = \arg(b)$ and $|p/q| = 1$). The plots show, that the ratios R_l approach unity for $a \approx \pm b\beta$. (Vanishing of R_l for $a \approx \pm b\beta$ follows also from the formula (19) if one takes into account that for $a \rightarrow b \pm \beta$ $\lambda_R \sim 1/(a \mp b\beta)$ whereas $|\mathcal{A}_R|^2 \sim |a \mp b\beta|^2$.) Therefore, for $a \approx \pm b$ the deviation from unity of R_μ is tiny while for R_τ it can be quite substantial.

The asymmetries $A_{\text{CP}}^{1,2}$ (13), (14) and the ratio R_l (15) are the observables in which the signal of CP violation (*i.e.* a non-zero phase δ_{CP}) vanishes for $a \approx \pm b\beta$. As the spin density matrix formalism will show, transverse polarization of the final state leptons is free from this problem. However, observables sensitive to the transverse polarization can experimentally be constructed only for τ 's which decay into hadrons. Thus, for $a \approx \pm b$ *only* the $\tau^+\tau^-$ channel provides an interesting opportunity to look for CP asymmetry in the leptonic $B_d^0(\bar{B}_d^0)$ decays.

The coefficients a and b ($\bar{a}$ and $\bar{b}$) are constrained by the experimental limit Eq. (1), which in the case $a \approx \pm b$ gives

$$|a| \approx |b| \lesssim 4.9 \times 10^{-9}. \quad (28)$$

In the minimal flavour violation scenario, the parameters a and b are also constrained indirectly⁶ by the limit (2). Taking $(m_\tau/m_\mu)(|V_{td}|/|V_{ts}|) \sim 4$ this constraint is satisfied if

$$|a| \approx |b| \lesssim 2 \times 10^{-9}. \quad (29)$$

⁶The limit imposed by the $B_s^0\text{-}\bar{B}_s^0$ mass difference [29, 32] can be avoided on account of a different $\tan\beta$ and $M_{A,H}$ dependence as compared to the Higgs penguin diagram contributions to $B_d^0(\bar{B}_d^0) \rightarrow l^+l^-$ decays [16].




Figure 1: The ratios R_μ and R_τ as functions of b/a for the phase $\delta_{\text{CP}} = -\frac{1}{2}\arg(\lambda_L) = 0.1$ (solid line), 0.3 (dashed), 0.5 (dotted) and 0.75 (dash-dotted). $R_l(-\delta_{\text{CP}}) = R_l^{-1}(\delta_{\text{CP}})$.

In our simulations in Section 5 we conservatively set $|a| = |b| \lesssim 10^{-9}$ and treat both scenarios simultaneously, as all what matters are the values of $|a| = |b|$ and the single CP violating phase δ_{CP} . In the MFV case $\delta_{\text{CP}} = \arg(\mu^* A_t^*)$ while in the NMFV case it is given by Eq. (24); in both cases the phase can be of order 1.

Finally let us notice that the SM as well as supersymmetric box and Z^0 penguin contributions spoil the exact equality $a = \pm b$. In addition, a finite difference of A^0 and H^0 masses also splits these two coefficients. It is therefore likely that $|a|$ and $|b|$ and the phases of a and b differ from each other by some 10÷15% even for $|a|$ and $|b|$ saturating the bounds (1), (2). Nevertheless, our simplified analysis will demonstrate that in supersymmetry there are good reasons to expect substantial CP violation in $B_d^0(\bar{B}_d^0) \rightarrow \tau^+\tau^-$ decays. Therefore in the following sections we present tools which can be used to search for these decays and look for CP violation.

4 Spin density matrix formalism

To see how the CP asymmetry in $B_d^0(\bar{B}_d^0) \rightarrow \tau^+\tau^-$ decays are reflected in realistic observables we use the TAUOLA τ -lepton decay library which allows us to simulate fully the effects of τ polarization. The exhaustive description of the method and numerical algorithm is given in papers [19] and [22]. The input to the TAUOLA Universal Interface [24] is the spin density matrix of the $\tau^+\tau^-$ system resulting from the decay of a neutral particle. In this section we collect the necessary formulae for this matrix for a $\tau^+\tau^-$ pair originating

from $B^0(\bar{B}^0)$.

The time dependent B^0 -meson mixing can easily be dealt with by introducing time dependent effective factors a_{eff} , b_{eff} , $\bar{a}_{\text{eff}}$ and $\bar{b}_{\text{eff}}$ defined as:

$$\begin{aligned} a_{\text{eff}}(t) &= a g_+(t) + \bar{a} \frac{q}{p} g_-(t) = a g_+(t) - a^* \frac{q}{p} g_-(t) , \\ b_{\text{eff}}(t) &= b g_+(t) + \bar{b} \frac{q}{p} g_-(t) = b g_+(t) + b^* \frac{q}{p} g_-(t) , \\ \bar{a}_{\text{eff}}(t) &= \bar{a} g_+(t) + a \frac{p}{q} g_-(t) = -a^* g_+(t) + a \frac{p}{q} g_-(t) , \\ \bar{b}_{\text{eff}}(t) &= \bar{b} g_+(t) + b \frac{p}{q} g_-(t) = b^* g_+(t) + b \frac{p}{q} g_-(t) , \end{aligned}$$

so that the instantaneous B^0 widths into left- or right-handed τ 's read

$$\Gamma(B_{\text{phys}}^0(t) \rightarrow \tau_L^+ \tau_L^-) = \frac{M_B}{16\pi} \beta |a_{\text{eff}}(t) + \beta b_{\text{eff}}(t)|^2 , \quad (30)$$

$$\Gamma(B_{\text{phys}}^0(t) \rightarrow \tau_R^+ \tau_R^-) = \frac{M_B}{16\pi} \beta |a_{\text{eff}}(t) - \beta b_{\text{eff}}(t)|^2 , \quad (31)$$

and those for $\bar{B}^0$ are given by similar formulae with $a_{\text{eff}}(t)$, $b_{\text{eff}}(t)$ replaced by $\bar{a}_{\text{eff}}(t)$, $\bar{b}_{\text{eff}}(t)$. CP is violated because in general $\bar{a}_{\text{eff}}(t) \neq -a_{\text{eff}}^*(t)$, and $\bar{b}_{\text{eff}}(t) \neq b_{\text{eff}}^*(t)$.

The spin weight for the $\tau^+ \tau^-$ pair originating from B^0 decay at time t is given by

$$wt = \frac{1}{4} \frac{T(s_1, p_1, s_2, p_2)}{T(0, p_1, 0, p_2)} , \quad (32)$$

where

$$T(s_1, p_1, s_2, p_2) = \text{Tr} \left[P(s_1) (\not{p}_1 + m_l) (b_{\text{eff}} + a_{\text{eff}} \gamma^5) P(s_2) (\not{p}_2 - m_l) (b_{\text{eff}}^* - a_{\text{eff}}^* \gamma^5) \right] ,$$

with $P(s) \equiv \frac{1}{2}(1 + \gamma^5 \not{s})$, and p_1, s_1 (p_2, s_2) are the momentum and spin four-vectors of the τ^- (τ^+) lepton, respectively. The τ -lepton spin four-vectors s_a in the B^0 ($\bar{B}^0$) rest frame are related to the spin three-vectors $\vec{\sigma}_a$ in the τ_a -lepton rest frames as follows⁷

$$\begin{aligned} s_1^0 &= \frac{M_B}{2m_l} \beta \sigma_1^z , & s_2^0 &= -\frac{M_B}{2m_l} \beta \sigma_2^z , \\ s_a^z &= \frac{M_B}{2m_l} \sigma_a^z , & s_a^{x,y} &= \sigma_a^{x,y} . \end{aligned} \quad (33)$$

Combining (32) with the τ -lepton decay matrix elements, the τ -lepton rest frame spin vectors $\vec{\sigma}_1$ and $\vec{\sigma}_2$ get replaced by the polarimetric vectors $\vec{h}_1$ and $\vec{h}_2$ which are determined solely by the dynamic of the considered τ decay process.

⁷In the rest frame of $B^0/\bar{B}^0$ the z -axis is aligned with τ^- momentum, exactly as in **TAUOLA Universal Interface**. However, here as a particle number 1 the τ^- lepton is taken (not τ^+), resulting in the transposition of the matrix $R_{\mu\nu}$ in the **TAUOLA Universal Interface**.

The spin weight for the complete event ($B \rightarrow \tau\tau \rightarrow \text{decay products}$) can be written in the form

$$WT = \frac{1}{4} \left(1 + \sum_{i=x,y,z} \sum_{j=x,y,z} R_{ij} h_1^i h_2^j + \sum_{i=x,y,z} R_{i0} h_1^i + \sum_{j=x,y,z} R_{0j} h_2^j \right). \quad (34)$$

Expanding (32) and comparing with (34) we find

$$\begin{aligned} R_{00} &= +1, & R_{x0} &= R_{y0} = R_{0x} = R_{0y} = 0, \\ R_{zz} &= -1, & R_{xz} &= R_{yz} = R_{zx} = R_{zy} = 0, \\ R_{0z} &= -R_{z0} = \frac{2\text{Re}(a_{\text{eff}} b_{\text{eff}}^*) \beta}{|b_{\text{eff}}|^2 \beta^2 + |a_{\text{eff}}|^2}, \\ R_{xy} &= -R_{yx} = -\frac{2\text{Im}(a_{\text{eff}} b_{\text{eff}}^*) \beta}{|b_{\text{eff}}|^2 \beta^2 + |a_{\text{eff}}|^2}, & R_{xx} &= R_{yy} = \frac{|b_{\text{eff}}|^2 \beta^2 - |a_{\text{eff}}|^2}{|b_{\text{eff}}|^2 \beta^2 + |a_{\text{eff}}|^2}, \end{aligned} \quad (35)$$

Eq. (34) with $R_{\mu\nu}$ replaced by $\bar{R}_{\mu\nu}$ computed as above but with $\bar{a}_{\text{eff}}(t)$ and $\bar{b}_{\text{eff}}(t)$ replacing $a_{\text{eff}}(t)$ and $b_{\text{eff}}(t)$, respectively, gives the spin weight for the events from $\bar{B}^0$ decays. CP violating effects in the $B^0 \rightarrow \tau^+ \tau^-$ and $\bar{B}^0 \rightarrow \tau^+ \tau^-$ decays are absent if $R_{0z} = \bar{R}_{z0}$, $R_{xy} = \bar{R}_{yx}$ and $R_{xx} = \bar{R}_{xx}$.

For simulations of the time integrated measurements, the time averaged matrix $\langle R_{\mu\nu} \rangle$ has to be used ⁸

$$\langle R_{\mu\nu} \rangle \equiv \frac{\int dt \Gamma(B_{\text{phys}}^0(t) \rightarrow \tau^+ \tau^-) R_{\mu\nu}(t)}{\int dt \Gamma(B_{\text{phys}}^0(t) \rightarrow \tau^+ \tau^-)}$$

and $\langle \bar{R}_{\mu\nu} \rangle$ given by a similar formula. The asymmetry (13) is then given by

$$A_{\text{CP}}^1 = \frac{(1 - \langle R_{z0} \rangle) \Gamma_{\text{int}} - (1 + \langle \bar{R}_{z0} \rangle) \bar{\Gamma}_{\text{int}}}{(1 - \langle R_{z0} \rangle) \Gamma_{\text{int}} + (1 + \langle \bar{R}_{z0} \rangle) \bar{\Gamma}_{\text{int}}}, \quad (36)$$

where

$$\Gamma_{\text{int}} = \int dt \Gamma(B_{\text{phys}}^0(t) \rightarrow \tau^+ \tau^-), \quad \bar{\Gamma}_{\text{int}} = \int dt \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow \tau^+ \tau^-).$$

$A_{\text{CP}}^2(t_1, t_2)$ defined in (14) is given by (36) reversing the signs in the brackets. It is also easy to check that for $|q/p| = 1$ and $a = \pm b$ one has $\langle R_{z0} \rangle \Gamma_{\text{int}} = -\langle \bar{R}_{z0} \rangle \bar{\Gamma}_{\text{int}}$ (*i.e.* $\text{Re}(a_{\text{eff}} b_{\text{eff}}^*) = -\text{Re}(\bar{a}_{\text{eff}} \bar{b}_{\text{eff}}^*)$). Then the two factors: $\langle R_{z0} \rangle \Gamma_{\text{int}}$ and $\langle \bar{R}_{z0} \rangle \bar{\Gamma}_{\text{int}}$ cancel out in the numerator of (36) and of the similar formula for $A_{\text{CP}}^2(t_1, t_2)$. As a result, nonzero asymmetries $A_{\text{CP}}^{1,2}$ are possible only if $\Gamma_{\text{int}} \neq \bar{\Gamma}_{\text{int}}$, which in view of the equalities $|a_{\text{eff}}(t)| = |\bar{b}_{\text{eff}}(t)|$, $|b_{\text{eff}}(t)| = |\bar{a}_{\text{eff}}(t)|$ requires β significantly different from 1. This again confirms our observation made in Section 2 that for $a \approx \pm b$ the asymmetry in the $\mu^+ \mu^-$ channel is suppressed.

⁸Simulations for time non-integrated measurements, with time-dependent decays explicitly generated are also possible.

Since in the limit $a = \pm b\beta$ the two integrated rates Γ_{int} and $\bar{\Gamma}_{\text{int}}$ are equal one gets $\langle R_{z0} \rangle = -\langle \bar{R}_{z0} \rangle$, which means that even for $\delta_{\text{CP}} \neq 0$ there can be no CP violating effects in the observables sensitive to the longitudinal polarization of τ 's (nor in R_τ). In contrast, it is easy to check by using the explicit analytical expressions, that in this limit the elements xx and xy of these matrices *need not* satisfy $\langle R_{xy} \rangle = -\langle \bar{R}_{xy} \rangle$ and $\langle R_{xx} \rangle = \langle \bar{R}_{xx} \rangle$. Hence, the observables sensitive to transverse τ polarization can reveal CP violation even if the observables introduced in Section 2 fail to signal it.

5 Results of the Monte Carlo simulations

As $B^0 \rightarrow \tau^+\tau^-$ decays have not yet been seen it might seem premature to study differential distributions in this channel, including τ polarization. Nevertheless, our analysis can serve as a good starting point for the future experimental work if indeed accumulated experimental samples would turn to be large enough. By providing a Monte Carlo tool useful in calculating *e.g.* detector responses, our study may also be considered as benchmarks for the simulations to be used in setting the upper limit for $B^0 \rightarrow \tau^-\tau^+$ branching ratios.

There are some similarities between this study and the one aiming at assessing measurability of the Higgs boson parity at future Linear Colliders [28]. The difference is that now the coefficients a and b (and also $\bar{a}$ and $\bar{b}$) in Eq. (3) can be complex, while in [28] they were taken to be purely imaginary and real, respectively. As a result, in the formula Eq. (34) terms linear in the polarimetric vector components also appear. Therefore we have extended the **TAUOLA Universal Interface** to include such a possibility as well. The algorithm for simulating B^0 decay into τ leptons is nearly identical to the one for H^0 decay. Changes necessary for implementation in the **Universal Interface** of the **TAUOLA** Monte Carlo library are limited to the replacement of Higgs identifier with the one for B_d^0 and $\bar{B}_d^0$ mesons, and the values of the spin density matrix with the ones computed in the preceding section.

The formulae given in the previous section are general. However, motivated by the two supersymmetric scenarios discussed in Section 3, in our numerical study we will show first the results obtained in the limit $a = \pm b$. Then we will discuss effects of a small departure from this relation.

We will limit ourselves to two observables, which are known to provide valuable and complementary information on the spin state of decaying τ lepton pairs.

$\pi^\pm$ energies: As the first observable we take the π^+ and π^- energy spectra in the decay channels $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ (or $\tau^- \rightarrow \pi^- \nu_\tau$). Since they reflect the longitudinal polarization of the individual $\tau^\pm$ leptons, the spectra are sensitive to R_{z0} and $\bar{R}_{0z}$ as can be inferred from the expression (34), *i.e.* they are sensitive to $\text{Re}(a_{\text{eff}} b_{\text{eff}}^*)$ as follows from (35). The CP violation is reflected in the difference between the energy spectrum of $\pi^-(\pi^+)$ originating from $B^0(\bar{B}^0)$ and the energy spectrum of $\pi^+(\pi^-)$ originating from $B^0(\bar{B}^0)$. This observable

was exploited before, for example in the $Z^0/\gamma^* \rightarrow \tau^+\tau^-$ study (for a review of the method and its extensions, see *e.g.* Ref. [35]). We will show energy spectra in the rest frame of the $B^0(\bar{B}^0)$ meson assuming that the reconstruction of the event kinematics in the BELLE and BABAR experiments is sufficiently good for that purpose, that is, that the momenta of τ decay products in the rest frame of $B^0(\bar{B}^0)$ can be reconstructed with the precision of a fraction of the τ mass.

Acoplanarity angle φ^* : As the second observable we use the acoplanarity angle φ^* between two planes spanned by the momenta of decay products of $\rho^\pm \rightarrow \pi^\pm\pi^0$ coming from decays of both τ leptons into $\rho\nu_\tau$ [28]. This quantity is sensitive to correlations between transverse components of τ -lepton spins (*i.e.* to the elements R_{xx} and R_{xy} which in turn probe $\text{Im}(a_{\text{eff}}b_{\text{eff}}^*)$, as can be seen from (35)). For the definition of the acoplanarity the orientation of decay planes and pion momenta has to be properly taken into account. The acoplanarity angle φ^* is defined with the help of two vectors $\mathbf{n}_\pm$ normal to the planes determined by the momenta of pions which originate from $\rho^\pm$ decays: $\mathbf{n}_\pm = \mathbf{p}_{\pi^\pm} \times \mathbf{p}_{\pi^0}$. If $\cos \xi = \frac{\mathbf{n}_+ \cdot \mathbf{n}_-}{|\mathbf{n}_+||\mathbf{n}_-|}$ then

$$\varphi^* = \begin{cases} \xi & \text{for } \text{sgn}(\mathbf{p}_{\pi^-} \cdot \mathbf{n}_+) < 0 \\ 2\pi - \xi & \text{for } \text{sgn}(\mathbf{p}_{\pi^-} \cdot \mathbf{n}_+) > 0 \end{cases} \quad (37)$$

making the full range of the variable $0 < \varphi^* < 2\pi$ of physical interest. Note that under CP, $\varphi^* \rightarrow 2\pi - \varphi^*$ since the condition $\text{sgn}(\mathbf{p}_{\pi^-} \cdot \mathbf{n}_+)$ is always evaluated from the orientation of π^- momentum with respect to the vector $\mathbf{n}_+$. In addition, we also have to sort events depending whether $y_1y_2 > 0$ or $y_1y_2 < 0$, where

$$y_1 = \frac{E_{\pi^+} - E_{\pi^0}}{E_{\pi^+} + E_{\pi^0}}, \quad y_2 = \frac{E_{\pi^-} - E_{\pi^0}}{E_{\pi^-} + E_{\pi^0}}, \quad (38)$$

since otherwise the spin correlations are washed out, as explained in Ref. [36]. The best would be to use in (38) the energies of $\pi^\pm$ and π^0 's in the rest frames of the corresponding $\tau^\pm$ leptons, but they are not directly measurable. Since in the $B^0(\bar{B}^0)$ rest frame the τ leptons are only mildly relativistic, the difference of pion energies in this frame and respective rest frames of $\tau^\pm$ should not be very important. The acoplanarity distribution is then evaluated in the rest frame of the $\rho^+\rho^-$ pair, but with the energies of $\pi^\pm$ and π^0 's in (38) taken in the rest frame of the $B^0(\bar{B}^0)$. The CP violation is reflected in the difference between the distributions of the acoplanarity angle φ^* measured in B^0 decays and the angle $2\pi - \varphi^*$ measured in $\bar{B}^0$ decays for the same signs of y_1y_2 .

Fig. 2 shows the pion energy spectra and the acoplanarity distributions assuming $|q/p| = 1$, $a = b = 10^{-9}$ and the CP violating phase $\delta_{\text{CP}} = 0.7$. For all plots the same number of 5×10^5 $\tau^+\tau^-$ events from B_d^0 and $\bar{B}_d^0$ decays has been generated with TAUOLA, although for the parameters chosen the ratio $R_\tau = 1.32$, see Fig. 1. In the upper left panel the energy spectra of π^- from B^0 decays (thick line with the slope proportional to $\langle R_{z0} \rangle$ [37]) and of π^+ from $\bar{B}^0$ (thin line; slope $\propto \langle \bar{R}_{0z} \rangle$) are shown, while in the lower left panel shown are the spectra of π^+ from B^0 decays (thick line; slope $\propto \langle R_{0z} \rangle$) and of π^- from $\bar{B}^0$ (thin line; slope $\propto \langle \bar{R}_{z0} \rangle$). The harder π^- energy spectrum from B_d^0 decays




Figure 2: *Results for the CP violating phase $\delta_{CP} = 0.7$ and $a = b$. Left panels: Single $\pi^\pm$ energy spectra in $B^0(\bar{B}^0) \rightarrow \tau^+\tau^-$, $\tau^\pm \rightarrow \pi^\pm \nu_\tau(\bar{\nu}_\tau)$. In the upper (lower) panel the thick line corresponds to the energy spectrum of π^- (of π^+) from B^0 decays and the thin line to the energy spectrum of π^+ (of π^-) from $\bar{B}^0$. Spectra are plotted in the rest frame of $B^0(\bar{B}^0)$. Right panels: acoplanarity distributions of the $\rho^+\rho^-$ decay products in $B^0(\bar{B}^0) \rightarrow \tau^+\tau^-$, $\tau^\pm \rightarrow \rho^\pm \nu_\tau(\bar{\nu}_\tau)$, $\rho^\pm \rightarrow \pi^\pm \pi^0$. The thick lines correspond to the acoplanarity angle φ^* measured in B^0 decays and the thin ones are for the angle $2\pi - \varphi^*$ measured in $\bar{B}^0$ decays. The acoplanarity angles are defined in the rest frame of the $\rho^+\rho^-$ pair. Events in the upper (lower) panel have $y_1y_2 > 0$ ($y_1y_2 < 0$).*

than π^+ from $\bar{B}_d^0$ (i.e. larger slope of the thick line) in the upper left panel indicates that $Br(B_d^0 \rightarrow \tau_R^+ \tau_R^-) > Br(\bar{B}_d^0 \rightarrow \tau_L^+ \tau_L^-)$, which is a clear signal of CP violation. In the acoplanarity plots (right panels) thick lines correspond to the distribution of φ^* measured in B^0 decays, and the thin lines to the distribution of $2\pi - \varphi^*$ measured in $\bar{B}^0$ decays; in the upper right panel $y_1y_2 > 0$, and $y_1y_2 < 0$ in the lower right one. The shapes of the thick and thin lines are described by the formulae

$$N_B(\varphi^*) = \text{const} - \text{sgn}(y_1y_2)A_R \cos(\varphi^* - \delta_R) \quad \text{for } B^0 \rightarrow \tau^+\tau^-$$

$$N_{\bar{B}}(\varphi^*) = \text{const} - \text{sgn}(y_1 y_2) A_{\bar{R}} \cos(2\pi - \varphi^* + \delta_{\bar{R}}) \quad \text{for } \bar{B}^0 \rightarrow \tau^+ \tau^-$$

where $A_R = (R_{xx}^2 + R_{xy}^2)^{1/2}$, $\sin \delta_R = R_{xy}/A_R$, $\cos \delta_R = R_{xx}/A_R$ and $A_{\bar{R}}$ and $\delta_{\bar{R}}$ are given by analogous formulae with R_{ij} replaced by the $\bar{R}_{ij}$. Different shapes of thick and thin lines seen in the right panels of Fig. 2 again indicate CP violation. In both energy and acoplanarity plots the CP violation is clearly seen and should be measurable even for small statistics.

Note also that if upper and lower plots are combined (*i.e.* no sorting according to the pion charge or sign of $y_1 y_2$ is made), all CP asymmetries are lost. Since the lower plots are simple reflections of the upper ones, in the following only plots corresponding to the upper panels of Fig. 2 are shown.




Figure 3: As in the upper panels of figure 2 but for $\delta_{\text{CP}} = 0.3$ and $a = b$.

Fig. 3 shows the corresponding distributions for the CP violating phase $\delta_{\text{CP}} = 0.3$ keeping still $a = b$ and $|q/p| = 1$. It is apparent that with decreasing $|\delta_{\text{CP}}|$ the signal of CP violation deteriorates (especially in the pion spectra) and the possibility of distinguishing pion spectra and acoplanarity distributions from B^0 and $\bar{B}^0$, and hence the CP violation, would require increasingly large statistics which may not be attainable at BELLE and BABAR without major upgrades.

As we discussed, the relation $a = \pm b$ is only approximate. Therefore, in Figure 4 (5) we show the $\pi^\pm$ spectra and acoplanarities for a greater (smaller) than b , but keeping as previously their phases equal (and assuming $|q/p| = 1$). We take $b = 0.8 a$ in Fig. 4 and $a = 0.8 b$ in Fig. 5. In both figures the single CP violating phase $\delta_{\text{CP}} = 0.7$.

Fig 4 shows that for $b = 0.8 a$ with the same value of δ_{CP} the CP violating effects in $\pi^\pm$ energy spectra are enhanced, while in the acoplanarities they are only slightly affected compared to the case $a = b$, c.f. the upper right panel of Fig. 2.

On the other hand, for a approaching $b\beta$ (for $B \rightarrow \tau^+ \tau^-$ decays $\beta \approx 0.74$) the effects of CP violation in the $\pi^\pm$ energy spectra disappear. This is clearly seen by comparing the

right panel of Fig. 5 with the upper left one of Fig. 2. This agrees with our observations following Eqs. (25) and (26) and with the discussion in Section 3. In contrast, the acoplanarities shown in the right panel of Fig. 5 clearly indicate the CP violation even for $a \approx b\beta$ confirming our discussion in Section 4. This clearly demonstrates the complementarity of the energy and acoplanarity distributions as a means to detect CP violation.




Figure 4: As in the upper panels of figure 2 but for $\delta_{CP} = 0.7$ and $b = 0.8a$.




Figure 5: As in the upper panels of figure 2 but for $\delta_{CP} = 0.7$ and $a = 0.8b$.

6 Conclusions

In this letter we have investigated possible signals of CP violation in the decays $B^0(\bar{B}^0) \rightarrow \tau^+\tau^-$. We have developed the necessary formalism and numerical tools allowing to apply

the **TAUOLA** τ -lepton decay library together with its **universal interface** to simulate fully the effects of the polarization of τ^+ and τ^- originating from such decays.

We have argued that in the interesting new physics scenario of supersymmetry with $\tan\beta \sim 40 \div 50$, in which the rates of $B_d^0(\bar{B}_d^0) \rightarrow \tau^+\tau^-$ decays are enhanced and could be detectable in the SLAC and KEK B -factories, the dependence of the CP asymmetries on the model parameters simplifies. Moreover, the CP violating phase needs not be small. In the non-minimal flavour violation case it can be of the same order as the phase of the V_{td} element of the CKM matrix. Therefore the CP asymmetries can be quite large as opposed to the $B^0(\bar{B}^0) \rightarrow \mu^+\mu^-$ decays in which they are kinematically suppressed.

By using Monte-Carlo simulations we have investigated the possible effects of CP violation in two realistic experimental observables and demonstrated that they might be detectable if the CP violating phase is reasonably large, *i.e.* $\mathcal{O}(1)$.

Since the decays $B^0(\bar{B}^0) \rightarrow \tau^+\tau^-$ have not been discovered yet, we have not discussed the statistics requirements nor attempted at including in our analysis any systematic or detector uncertainties. It is clear that once these decays are discovered, other τ decay channels than the ones investigated here can be analysed jointly to give additional information on the polarization of τ 's. Our numerical tools are prepared for that. The tools can also be applied to determine the upper limits on the branching fraction of the $B^0(\bar{B}^0) \rightarrow \tau^+\tau^-$ decays by the *BABAR* and *BELLE* collaborations.

As a final remark, we point that our analysis can be taken over to Higgs boson production at linear colliders with its subsequent decay to τ pairs, which as yet has not been simulated in connection with the complex scalar and pseudoscalar couplings.

Acknowledgements

The authors would like to thank P. Ball, A. Martin, M. Misiak, H. Pałka and J. Stirling for useful discussions. P.H.Ch. thanks the CERN Theory Group for hospitality during the initial stage of this research.

Work supported by the Polish State Committee for Scientific Research Grants 2 P03B 040 24 for years 2003-2005 (P.H.Ch. and J.K.), 1 P03 091 27 for years 2004-2006 (Z.W.), 1 P03B 009 27 for years 2004-2005 (M.W.), and by the EC Contracts HPRN-CT-2000-00148 (P.H.Ch.) and HPRN-CT-2000-00149 (J.K). In addition, M.W. acknowledges the Maria Skłodowska-Curie Fellowship granted by the the European Community in the framework of the Human Potential Programme under contract HPMD-CT-2001-00105 (*"Multi-particle production and higher order correction"*).

References

- [1] K. Abe *et al.* [BELLE Collaboration], arXiv:hep-ex/0409049.
A. Hocker [BABAR Collaboration], arXiv:hep-ex/0410069.

- [2] W. Skiba and J. Kalinowski, Nucl. Phys. B **404** (1993) 3. H. E. Logan and U. Nierste, Nucl. Phys. B **586** (2000) 39 [arXiv:hep-ph/0004139].
- [3] C. Hamzaoui, M. Pospelov and M. Toharia, *Phys. Rev.* **D59** (1999) 095005.
- [4] K. S. Babu and C. F. Kolda, Phys. Rev. Lett. **84** (2000) 228 [arXiv:hep-ph/9909476].
- [5] P. H. Chankowski and L. Slawianowska, Phys. Rev. D **63** (2001) 054012 [arXiv:hep-ph/0008046].
- [6] G. Isidori and A. Retico, JHEP **0111** (2001) 001 [arXiv:hep-ph/0110121].
- [7] C. S. Huang and W. Liao, Phys. Lett. B **525** (2002) 107 [arXiv:hep-ph/0011089].
- [8] A. Dedes and A. Pilaftsis, Phys. Rev. D **67** (2003) 015012 [arXiv:hep-ph/0209306].
- [9] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0408095.
- [10] M. C. Chang *et al.* [BELLE Collaboration], Phys. Rev. D **68** (2003) 111101 [arXiv:hep-ex/0309069].
- [11] D. Acosta *et al.* [CDF Collaboration], Phys. Rev. Lett. **93** (2004) 032001 [arXiv:hep-ex/0403032].
- [12] M. Herndon, talk given at 32nd *Int. Conf. on High Energy Physics* (ICHEP 2004), Beijing, China, August 2004.
- [13] V. M. Abazov *et al.* [D0 Collaboration], arXiv:hep-ex/0410039.
- [14] see e.g. A. J. Buras, in *Probing the Standard Model of Electroweak Interactions*, F. David and R. Gupta eds., Elsevier Science B.V., 1998 [arXiv:hep-ph/9806471].
- [15] P. Ball *et al.*, in *Standard Model Physics (and more) at the LHC*, G. Altarelli and M. L. Mangano eds., CERN, 2000 [arXiv:hep-ph/0003238].
- [16] P. H. Chankowski and L. Slawianowska, Eur. Phys. J. C **33**, 123 (2004) [arXiv:hep-ph/0308032].
- [17] S. Hashimoto, talk given at 32nd *Int. Conf. on High Energy Physics* (ICHEP 2004), Beijing, China, August 2004, arXiv:hep-ph/0411126.
- [18] C. S. Huang and W. Liao, arXiv:hep-ph/0211131; Phys. Lett. B **538** (2002) 301 [arXiv:hep-ph/0201121].
- [19] S. Jadach, J. H. Kuhn and Z. Was, Comput. Phys. Commun. **64** (1990) 275.
- [20] M. Jezabek, Z. Was, S. Jadach and J. H. Kuhn, Comput. Phys. Commun. **70** (1992) 69.
- [21] S. Jadach, Z. Was, R. Decker and J. H. Kuhn, Comput. Phys. Commun. **76** (1993) 361.

- [22] Z. Was and M. Worek, *Acta Phys. Polon. B* **33** (2002) 1875 [arXiv:hep-ph/0202007].
- [23] T. Pierzchala, E. Richter-Was, Z. Was and M. Worek, *Acta Phys. Polon. B* **32** (2001) 1277 [arXiv:hep-ph/0101311].
- [24] P. Golonka *et al.*, arXiv:hep-ph/0312240.
- [25] G.C. Branco, F. Lavoura and J.P. Silva, *CP violation*, Oxford University Press, Oxford, UK 1999; I.I.Y. Bigi and A.I. Sanda, *CP violation*, Cambridge University Press, Cambridge, UK 2000.
- [26] P. H. Chankowski and L. Slawianowska, *Acta Phys. Polon. B* **32** (2001) 1895.
- [27] Y. S. Tsai, *Phys. Rev. D* **4** (1971) 2821 [Erratum-ibid. *D* **13** (1976) 771]. H. K. Kuhn and F. Wagner, *Nucl. Phys. B* **236** (1984) 16. B. K. Bullock, K. Hagiwara and A. D. Martin, *Phys. Rev. Lett.* **67** (1991) 3055, *Nucl. Phys. B* **395** (1993) 499.
- [28] K. Desch, A. Imhof, Z. Was and M. Worek, *Phys. Lett. B* **579** (2004) 157 [arXiv:hep-ph/0307331].
- [29] A. J. Buras, P. H. Chankowski, J. Rosiek and L. Slawianowska, *Phys. Lett. B* **546** (2002) 96 [arXiv:hep-ph/0207241]; *Nucl. Phys. B* **659** (2003) 3 [arXiv:hep-ph/0210145].
- [30] G. Isidori and A. Retico, *JHEP* **0209** (2002) 063 [arXiv:hep-ph/0208159].
- [31] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, *Nucl. Phys. B* **477** (1996) 321 [arXiv:hep-ph/9604387]. M. Misiak, S. Pokorski and J. Rosiek, *Adv. Ser. Direct. High Energy Phys.* **15** (1998) 795 [arXiv:hep-ph/9703442]. D. Becirevic *et al.*, *Nucl. Phys. B* **634** (2002) 105 [arXiv:hep-ph/0112303].
- [32] A. J. Buras, P. H. Chankowski, J. Rosiek and L. Slawianowska, *Nucl. Phys. B* **619** (2001) 434 [arXiv:hep-ph/0107048].
- [33] P. H. Chankowski and J. Rosiek, *Acta Phys. Polon. B* **33** (2002) 2329 [arXiv:hep-ph/0207242].
- [34] A. Dedes, *Mod. Phys. Lett. A* **18** (2003) 2627 [arXiv:hep-ph/0309233].
- [35] A. Heister *et al.* [ALEPH Collaboration], *Eur. Phys. J. C* **20** (2001) 401 [arXiv:hep-ex/0104038].
- [36] G. R. Bower, T. Pierzchala, Z. Was and M. Worek, *Phys. Lett. B* **543** (2002) 227 [arXiv:hep-ph/0204292].
- [37] For a pedagogical presentation, see P. H. Eberhard *et al.*, CERN-EP-89-129, in *Proceedings of the Workshop on Z Physics at LEP*, G. Altarelli, R. Kleiss and V. Verzegnassi eds., Geneva, Switzerland, 1989.